

PRESCRIBING INFORMATION


RELENZA[®]

(zanamivir for inhalation)

For Oral Inhalation Only
For Use with the DISKHALER[®] Inhalation Device

DESCRIPTION

The active component of RELENZA is zanamivir. The chemical name of zanamivir is 5-(acetylamino)-4-[(aminoiminomethyl)-amino]-2,6-anhydro-3,4,5-trideoxy-D-glycero-D-galactonon-2-enonic acid. It has a molecular formula of C₁₂H₂₀N₄O₇ and a molecular weight of 332.3. It has the following structural formula:



Zanamivir is a white to off-white powder with a solubility of approximately 18 mg/mL in water at 20°C.

RELENZA is for administration to the respiratory tract by oral inhalation only. Each RELENZA ROTADISK[®] contains 4 regularly spaced double-foil blisters with each blister containing a powder mixture of 5 mg of zanamivir and 20 mg of lactose (which contains milk proteins). The contents of each blister are inhaled using a specially designed breath-activated plastic device for inhaling powder called the DISKHALER. After a RELENZA ROTADISK is loaded into the DISKHALER, a blister that contains medication is pierced and the zanamivir is dispersed into the air stream created when the patient inhales through the mouthpiece. The amount of drug delivered to the respiratory tract will depend on patient factors such as inspiratory flow. Under standardized in vitro testing, RELENZA ROTADISK delivers 4 mg of zanamivir from the DISKHALER device when tested at a pressure drop of 3 kPa (corresponding to a flow rate of about 62 to 65 L/min) for 3 seconds. In a study of 5 adult and 5 adolescent patients with obstructive airway diseases, the combined peak inspiratory flow rates (PIFR) ranged from 66 to 140 L/min. In a separate study of 16 pediatric patients, PIFR results were more variable; 4 did not achieve measurable flow rates, and PIFR for measurable inhalations by 12 children ranged from 30.5 to 122.4 L/min. Only 1 of 4 children under age 8 had a measurable flow rate (see CLINICAL PHARMACOLOGY: Pediatric Patients, INDICATIONS AND USAGE: Description of Clinical Studies, and PRECAUTIONS: Pediatric Use).

MICROBIOLOGY

Mechanism of Action: The mechanism of action of zanamivir is via inhibition of influenza virus neuraminidase with the possibility of alteration of virus particle aggregation and release.

Antiviral Activity: The antiviral activity of zanamivir against laboratory and clinical isolates of influenza virus was determined in cell culture assays. The concentrations of zanamivir required for inhibition of influenza virus were highly variable depending on the assay method used and virus isolate tested. The 50% and 90% effective concentrations (EC_{50} and EC_{90}) of zanamivir were in the range of 0.005 to 16.0 μ M and 0.05 to >100 μ M, respectively

(1 μ M = 0.33 mcg/mL). The relationship between the in vitro inhibition of influenza virus by zanamivir and the inhibition of influenza virus replication in humans has not been established.

Resistance: Influenza viruses with reduced susceptibility to zanamivir have been recovered in vitro by multiple passages of the virus in the presence of increasing concentrations of the drug. Genetic analysis of these viruses showed that the reduced susceptibility in vitro to zanamivir is associated with mutations that result in amino acid changes in the viral neuraminidase or viral hemagglutinin or both. Resistance mutations selected in vitro which result in neuraminidase amino acid substitutions include E119G/A/D and R292K. Mutations selected in vitro in hemagglutinin include: K68R, G75E, E114K, N145S, S165N, S186F, N199S, and K222T.

In an immunocompromised patient infected with influenza B virus, a variant virus emerged after treatment with an investigational nebulized solution of zanamivir for 2 weeks. Analysis of this variant showed a hemagglutinin mutation (T198I) which resulted in a reduced affinity for human cell receptors, and a substitution in the neuraminidase active site (R152K) which reduced the enzyme's activity to zanamivir by 1,000-fold. Insufficient information is available to characterize the risk of emergence of zanamivir resistance in clinical use.

Cross-Resistance: Cross-resistance has been observed between some zanamivir-resistant and some oseltamivir-resistant influenza virus mutants generated in vitro. However, some of the in vitro zanamivir-induced resistance mutations, E119G/A/D and R292K, occurred at the same neuraminidase amino acid positions as in the clinical isolates resistant to oseltamivir, E119V and R292K. No studies have been performed to assess risk of emergence of cross-resistance during clinical use.

Influenza Vaccine Interaction Study: An interaction study (n = 138) was conducted to evaluate the effects of zanamivir (10 mg once daily) on the serological response to a single dose of trivalent inactivated influenza vaccine, as measured by hemagglutination inhibition titers. There was no clear difference in hemagglutination inhibition antibody titers at 2 weeks and 4 weeks after vaccine administration between zanamivir and placebo recipients.

Influenza Challenge Studies: Antiviral activity of zanamivir was supported for infection with influenza A virus, and to a more limited extent for infection with influenza B virus, by Phase 1 studies in volunteers who received intranasal inoculations of challenge strains of influenza virus, and received an intranasal formulation of zanamivir or placebo starting before or shortly after viral inoculation.

CLINICAL PHARMACOLOGY

Pharmacokinetics: Absorption and Bioavailability: Pharmacokinetic studies of orally inhaled zanamivir indicate that approximately 4% to 17% of the inhaled dose is systemically absorbed. The peak serum concentrations ranged from 17 to 142 ng/mL within 1 to 2 hours following a 10-mg dose. The area under the serum concentration versus time curve (AUC_{∞}) ranged from 111 to 1,364 ng•hr/mL.

Distribution: Zanamivir has limited plasma protein binding (<10%).

Metabolism: Zanamivir is renally excreted as unchanged drug. No metabolites have been detected in humans.

Elimination: The serum half-life of zanamivir following administration by oral inhalation ranges from 2.5 to 5.1 hours. It is excreted unchanged in the urine with excretion of a single dose completed within 24 hours. Total clearance ranges from 2.5 to 10.9 L/hr. Unabsorbed drug is excreted in the feces.

Special Populations: Impaired Hepatic Function: The pharmacokinetics of zanamivir have not been studied in patients with impaired hepatic function.

Impaired Renal Function: Systemic exposure is limited after inhalation (see Absorption and Bioavailability). After a single intravenous dose of 4 mg or 2 mg of zanamivir in volunteers with mild/moderate or severe renal impairment, respectively, significant decreases in renal clearance (and hence total clearance: normals 5.3 L/hr, mild/moderate 2.7 L/hr, and severe 0.8 L/hr; median values) and significant increases in half-life (normals 3.1 hr, mild/moderate 4.7 hr, and severe 18.5 hr; median values) and systemic exposure were observed. Safety and efficacy have not been documented in the presence of severe renal insufficiency.

Pediatric Patients: The pharmacokinetics of zanamivir were evaluated in pediatric patients with signs and symptoms of respiratory illness. Sixteen patients, 6 to 12 years of age, received a single dose of 10-mg zanamivir dry powder via DISKHALER. Five patients had either undetectable zanamivir serum concentrations or had low drug concentrations (8.32 to 10.38 ng/mL) that were not detectable after 1.5 hours. Eleven patients had C_{max} median values of 43 ng/mL (range 15 to 74) and AUC_{∞} median values of 167 ng•hr/mL (range 58 to 279). Low or undetectable serum concentrations were related to lack of measurable PIFR in individual patients (see DESCRIPTION, INDICATIONS AND USAGE: Description of Clinical Studies, and PRECAUTIONS: Pediatric Use).

Geriatric Patients: The pharmacokinetics of zanamivir have not been studied in patients over 65 years of age (see PRECAUTIONS: Geriatric Use).

Gender, Race, and Weight: In a population pharmacokinetic analysis in patient studies, no clinically significant differences in serum concentrations and/or pharmacokinetic parameters (V/F , CL/F , k_a , AUC_{0-3} , C_{max} , T_{max} , CL_r , and % excreted in urine) were observed when demographic variables (gender, age, race, and weight) and indices of infection (laboratory evidence of infection, overall symptoms, symptoms of upper respiratory illness, and viral titers) were considered. There were no significant correlations between measures of systemic exposure and safety parameters.

Drug Interactions: No clinically significant pharmacokinetic drug interactions are predicted based on data from in vitro studies.

Zanamivir is not a substrate nor does it affect cytochrome P450 (CYP) isoenzymes (CYP1A1/2, 2A6, 2C9, 2C18, 2D6, 2E1, and 3A4) in human liver microsomes.

INDICATIONS AND USAGE

Treatment of Influenza: RELENZA is indicated for treatment of uncomplicated acute illness due to influenza A and B virus in adults and pediatric patients 7 years of age and older who have been symptomatic for no more than 2 days (see Description of Clinical Studies and PRECAUTIONS).

Prophylaxis of Influenza: RELENZA is indicated in adults and pediatric patients 5 years of age and older for prophylaxis of influenza.

Important Information on Use of RELENZA:

- RELENZA is not recommended for treatment or prophylaxis of influenza in individuals with underlying airways disease (such as asthma or chronic obstructive pulmonary disease [see WARNINGS]) due to risk of serious bronchospasm.
- RELENZA has not been proven effective for treatment of influenza in individuals with underlying airways disease.
- RELENZA has not been proven effective for prophylaxis of influenza in the nursing home setting.
- RELENZA is not a substitute for early vaccination on an annual basis as recommended by the Centers for Disease Control's Immunization Practices Advisory Committee.

Description of Clinical Studies: *Treatment of Influenza: Adults and Adolescents:*

The efficacy of RELENZA 10 mg inhaled twice daily for 5 days in the treatment of influenza has been evaluated in placebo-controlled studies conducted in North America, the Southern Hemisphere, and Europe during their respective influenza seasons. The magnitude of treatment effect varied between studies, with possible relationships to population-related factors including amount of symptomatic relief medication used.

Populations Studied: The principal Phase 3 studies enrolled 1,588 patients ages 12 years and older (median age 34 years, 49% male, 91% Caucasian), with uncomplicated influenza-like illness within 2 days of symptom onset. Influenza was confirmed by culture, hemagglutination inhibition antibodies, or investigational direct tests. Of 1,164 patients with confirmed influenza, 89% had influenza A and 11% had influenza B. These studies served as the principal basis for efficacy evaluation, with more limited Phase 2 studies providing supporting information where necessary. Following randomization to either zanamivir or placebo (inhaled lactose vehicle), all patients received instruction and supervision by a healthcare professional for the initial dose.

Principal Results: The definition of time to improvement in major symptoms of influenza included no fever and self-assessment of “none” or “mild” for headache, myalgia, cough, and sore throat. A Phase 2 and a Phase 3 study conducted in North America (total of over

600 influenza-positive patients) suggested up to one day of shortening of median time to this defined improvement in symptoms in patients receiving zanamivir compared to placebo, although statistical significance was not reached in either of these studies. In a study conducted in the Southern Hemisphere (321 influenza-positive patients), a 1.5-day difference in median time to symptom improvement was observed. Additional evidence of efficacy was provided by the European study.

Other Findings: There was no consistent difference in treatment effect in patients with influenza A compared to influenza B; however, these trials enrolled smaller numbers of patients with influenza B and thus provided less evidence in support of efficacy in influenza B.

In general, patients with lower temperature (e.g., 38.2°C or less) or investigator-rated as having less severe symptoms at entry derived less benefit from therapy.

No consistent treatment effect was demonstrated in patients with underlying chronic medical conditions, including respiratory or cardiovascular disease (see WARNINGS and PRECAUTIONS).

No consistent differences in rate of development of complications were observed between treatment groups.

Some fluctuation of symptoms was observed after the primary study endpoint in both treatment groups.

Pediatric Patients: The efficacy of RELENZA 10 mg inhaled twice daily for 5 days in the treatment of influenza in pediatric patients has been evaluated in a placebo-controlled study conducted in North America and Europe, enrolling 471 patients, ages 5 to 12 years (55% male, 90% Caucasian), within 36 hours of symptom onset. Of 346 patients with confirmed influenza, 65% had influenza A and 35% had influenza B. The definition of time to improvement included no fever and parental assessment of no or mild cough and absent/minimal muscle and joint aches or pains, sore throat, chills/feverishness, and headache. Median time to symptom improvement was one day shorter in patients receiving zanamivir compared with placebo. No consistent differences in rate of development of complications were observed between treatment groups. Some fluctuation of symptoms was observed after the primary study endpoint in both treatment groups.

Although this study was designed to enroll children ages 5 to 12 years, the product is indicated only for children 7 years of age and older. This evaluation is based on the combination of lower estimates of treatment effect in 5- and 6-year-olds compared with the overall study population, and evidence of inadequate inhalation through the DISKHALER in a pharmacokinetic study (see DESCRIPTION, CLINICAL PHARMACOLOGY: Pediatric Patients, and PRECAUTIONS: Pediatric Use).

Prophylaxis of Influenza: The efficacy of RELENZA in preventing naturally occurring influenza illness has been demonstrated in 2 post-exposure prophylaxis studies in households and 2 seasonal prophylaxis studies during community outbreaks of influenza. The primary efficacy endpoint in these studies was the incidence of symptomatic, laboratory-confirmed influenza, defined as the presence of 2 or more of the following symptoms: oral temperature

≥100°F/37.8°C or feverishness, cough, headache, sore throat, and myalgia; and laboratory confirmation of influenza A or B by culture, PCR, or seroconversion (defined as a 4-fold increase in convalescent antibody titer from baseline).

Two studies assessed post-exposure prophylaxis in household contacts of an index case. Within 1.5 days of onset of symptoms in an index case, each household (including all family members ≥5 years of age) was randomized to RELENZA 10 mg inhaled once daily or placebo inhaled once daily for 10 days. In the first study only, each index case was randomized to RELENZA 10 mg inhaled twice daily for 5 days or inhaled placebo twice daily for 5 days. In this study, the proportion of households with at least 1 new case of symptomatic laboratory-confirmed influenza was reduced from 19.0% (32 of 168 households) for the placebo group to 4.1% (7 of 169 households) for the group receiving RELENZA.

In the second study, index cases were not treated. The incidence of symptomatic laboratory-confirmed influenza was reduced from 19.0% (46 of 242 households) for the placebo group to 4.1% (10 of 245 households) for the group receiving RELENZA.

Two seasonal prophylaxis studies assessed RELENZA 10 mg inhaled once daily versus placebo inhaled once daily for 28 days during community outbreaks. The first study enrolled subjects 18 years of age or greater (mean age 29 years) from two university communities. The majority of subjects were unvaccinated (86%). In this study, the incidence of symptomatic laboratory-confirmed influenza was reduced from 6.1% (34 of 554) for the placebo group to 2.0% (11 of 553) for the group receiving RELENZA.

The second seasonal prophylaxis study enrolled subjects 12 to 94 years of age (mean age 60 years) with 56% of them older than 65 years of age. Sixty-seven percent of the subjects were vaccinated. In this study, the incidence of symptomatic laboratory-confirmed influenza was reduced from 1.4% (23 of 1,685) for the placebo group to 0.2% (4 of 1,678) for the group receiving RELENZA.

CONTRAINDICATIONS

RELENZA is contraindicated in patients with a known hypersensitivity to any component of the formulation (see DESCRIPTION).

WARNINGS

RELENZA IS NOT RECOMMENDED FOR TREATMENT OR PROPHYLAXIS OF INFLUENZA IN INDIVIDUALS WITH UNDERLYING AIRWAYS DISEASE (SUCH AS ASTHMA OR CHRONIC OBSTRUCTIVE PULMONARY DISEASE) (see INDICATIONS AND USAGE).

Serious cases of bronchospasm, including fatalities, have been reported during treatment with RELENZA in patients with and without underlying airways disease. Many of these cases were reported during postmarketing and causality was difficult to assess.

RELENZA SHOULD BE DISCONTINUED IN ANY PATIENT WHO DEVELOPS BRONCHOSPASM OR DECLINE IN RESPIRATORY FUNCTION; immediate treatment and hospitalization may be required. Some patients without prior pulmonary

disease may also have respiratory abnormalities from acute respiratory infection that could resemble adverse drug reactions or increase patient vulnerability to adverse drug reactions.

Bronchospasm was documented following administration of zanamivir in 1 of 13 patients with mild or moderate asthma (but without acute influenza-like illness) in a Phase 1 study. In interim results from an ongoing treatment study in patients with acute influenza-like illness superimposed on underlying asthma or chronic obstructive pulmonary disease, more patients on zanamivir than on placebo experienced greater than 20% decline in FEV₁ or peak expiratory flow rate.

If treatment with RELENZA is considered for a patient with underlying airways disease, the potential risks and benefits should be carefully weighed. If a decision is made to prescribe RELENZA for such a patient, this should be done only under conditions of careful monitoring of respiratory function, close observation, and appropriate supportive care including availability of fast-acting bronchodilators.

PRECAUTIONS

General: Patients should be instructed in the use of the delivery system. Instructions should include a demonstration whenever possible. Patients should read and follow carefully the Patient Instructions for Use accompanying the product. Effective and safe use of RELENZA requires proper use of the DISKHALER to inhale the drug.

There is no evidence for efficacy of zanamivir in any illness caused by agents other than influenza virus A and B.

No data are available to support safety or efficacy in patients who begin treatment after 48 hours of symptoms.

Safety and efficacy of repeated treatment courses have not been studied.

Allergic Reactions: Allergic-like reactions, including oropharyngeal edema, serious skin rashes, and anaphylaxis have been reported in post-marketing experience with RELENZA. RELENZA should be stopped and appropriate treatment instituted if an allergic reaction occurs or is suspected.

Bacterial Infections: Serious bacterial infections may begin with influenza-like symptoms or may coexist with or occur as complications during the course of influenza. RELENZA has not been shown to prevent such complications.

Prevention of Influenza: Use of zanamivir should not affect the evaluation of individuals for annual influenza vaccination in accordance with guidelines of the Centers for Disease Control and Prevention Advisory Committee on Immunization Practices.

Limitations of Populations Studied: Safety and efficacy have not been demonstrated in patients with high-risk underlying medical conditions (see INDICATIONS AND USAGE: Description of Clinical Studies, and WARNINGS). No information is available regarding treatment of influenza in patients with any medical condition sufficiently severe or unstable to be considered at imminent risk of requiring inpatient management.

Information for Patients: Patients should be instructed in use of the delivery system. Instructions should include a demonstration whenever possible.

For the proper use of RELENZA, the patient should read and follow carefully the accompanying Patient Instructions for Use.

Patients should be advised that the use of RELENZA for treatment of influenza has not been shown to reduce the risk of transmission of influenza to others.

Patients should be advised of the risk of bronchospasm, especially in the setting of underlying airways disease, and should stop RELENZA and contact their physician if they experience increased respiratory symptoms during treatment such as worsening wheezing, shortness of breath, or other signs or symptoms of bronchospasm (see WARNINGS). If a decision is made to prescribe RELENZA for a patient with asthma or chronic obstructive pulmonary disease, the patient should be made aware of the risks and should have a fast-acting bronchodilator available. Patients scheduled to take inhaled bronchodilators at the same time as RELENZA should be advised to use their bronchodilators before taking RELENZA.

Drug Interactions: No clinically significant pharmacokinetic drug interactions are predicted based on data from in vitro studies.

Carcinogenesis, Mutagenesis, and Impairment of Fertility: Carcinogenesis: In 2-year carcinogenicity studies conducted in rats and mice using a powder formulation administered through inhalation, zanamivir induced no statistically significant increases in tumors over controls. The maximum daily exposures in rats and mice were approximately 23 to 25 and 20 to 22 times, respectively, greater than those in humans at the proposed clinical dose based on AUC comparisons.

Mutagenesis: Zanamivir was not mutagenic in in vitro and in vivo genotoxicity assays which included bacterial mutation assays in *S. typhimurium* and *E. coli*, mammalian mutation assays in mouse lymphoma, chromosomal aberration assays in human peripheral blood lymphocytes, and the in vivo mouse bone marrow micronucleus assay.

Impairment of Fertility: The effects of zanamivir on fertility and general reproductive performance were investigated in male (dosed for 10 weeks prior to mating, and throughout mating, gestation/lactation, and shortly after weaning) and female rats (dosed for 3 weeks prior to mating through day 19 of pregnancy, or day 21 post partum) at IV doses 1, 9, and 90 mg/kg/day. Zanamivir did not impair mating or fertility of male or female rats, and did not affect the sperm of treated male rats. The reproductive performance of the F1 generation born to female rats given zanamivir was not affected. Based on a subchronic study in rats at a 90-mg/kg/day IV dose, AUC values ranged between 142 and 199 mcg•hr/mL (>300 times the human exposure at the proposed clinical dose).

Pregnancy: Pregnancy Category C. Embryo/fetal development studies were conducted in rats (dosed from days 6 to 15 of pregnancy) and rabbits (dosed from days 7 to 19 of pregnancy) using the same IV doses. Pre- and post-natal developmental studies were performed in rats (dosed from day 16 of pregnancy until litter day 21 to 23). In all studies, intravenous (1, 9, and 90 mg/kg/day)

instead of the inhalational route of drug administration was used. No malformations, maternal toxicity, or embryotoxicity were observed in pregnant rats or rabbits and their fetuses. Because of insufficient blood sampling timepoints in both rat and rabbit reproductive toxicity studies, AUC values were not available. However, in a subchronic study in rats at the 90-mg/kg/day IV dose, the AUC values were greater than 300 times the human exposure at the proposed clinical dose.

An additional embryo/fetal study, in a different strain of rat, was conducted using subcutaneous administration of zanamivir, 3 times daily, at doses of 1, 9, or 80 mg/kg during days 7 to 17 of pregnancy. There was an increase in the incidence rates of a variety of minor skeleton alterations and variants in the exposed offspring in this study. Based on AUC measurements, the high dose in the study produced an exposure greater than 1,000 times the human exposure at the proposed clinical dose. However, the individual incidence rate of each skeletal alteration or variant, in most instances, remained within the background rates of the historical occurrence in the strain studied.

Zanamivir has been shown to cross the placenta in rats and rabbits. In these animals, fetal blood concentrations of zanamivir were significantly lower than zanamivir concentrations in the maternal blood.

There are no adequate and well-controlled studies of zanamivir in pregnant women. Zanamivir should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Nursing Mothers: Studies in rats have demonstrated that zanamivir is excreted in milk. However, nursing mothers should be instructed that it is not known whether zanamivir is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when RELENZA is administered to a nursing mother.

Pediatric Use: Safety and effectiveness of RELENZA for treatment of influenza have not been assessed in pediatric patients less than 7 years of age.

The safety and effectiveness of RELENZA have been studied in a Phase 3 treatment study in pediatric patients, where 471 children 5 to 12 years of age received zanamivir or placebo (see INDICATIONS AND USAGE: Description of Clinical Studies, ADVERSE REACTIONS, and DOSAGE AND ADMINISTRATION). In a Phase 1 study of 16 children ages 6 to 12 years with signs and symptoms of respiratory disease, 4 did not produce a measurable peak inspiratory flow rate (PIFR) through the DISKHALER (3 with no adequate inhalation on request, 1 with missing data), 9 had measurable PIFR on each of 2 inhalations, and 3 achieved measurable PIFR on only 1 of 2 inhalations. Neither of two 6-year-olds and one of two 7-year-olds produced measurable PIFR. Overall, 8 of the 16 children (including all those under 8 years old) either did not produce measurable inspiratory flow through the DISKHALER or produced peak inspiratory flow rates below the 60 L/min considered optimal for the device under standardized in vitro testing; lack of measurable flow rate was related to low or undetectable serum concentrations (see DESCRIPTION, CLINICAL PHARMACOLOGY: Pediatric Patients, and INDICATIONS AND USAGE: Description of Clinical Studies). Prescribers should carefully evaluate the ability of

young children to use the delivery system if prescription of RELENZA is considered. When RELENZA is prescribed for children, it should be used only under adult supervision and with attention to proper use of the delivery system.

Adolescents were included in the three principal Phase 3 adult treatment studies. In these studies, 67 patients were 12 to 16 years of age. No definite differences in safety and efficacy were observed between these adolescent patients and young adults.

In addition, the safety and effectiveness of RELENZA for prophylaxis of influenza have been studied in four Phase 3 studies where 273 children 5 to 11 years of age and 239 adolescents 12 to 16 years of age received RELENZA. No differences in safety and effectiveness were observed between pediatric and adult subjects.

Geriatric Use: Of the total number of patients in 6 clinical studies of RELENZA for treatment of influenza, 59 were 65 and over, while 24 were 75 and over. Of the total number of patients in 4 clinical studies of RELENZA for prophylaxis of influenza in households and community settings, 954 were 65 and over, while 347 were 75 and over. No overall differences in safety or effectiveness were observed between these subjects and younger patients, and other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

In 2 additional studies of RELENZA for prophylaxis of influenza in the nursing home setting, efficacy was not demonstrated (see INDICATIONS AND USAGE). Elderly subjects may need assistance with use of the device.

ADVERSE REACTIONS

See **WARNINGS and PRECAUTIONS for information about risk of serious adverse events such as bronchospasm and allergic-like reactions, and for safety information in patients with underlying airways disease.**

Because the placebo consisted of inhaled lactose powder, which is also the vehicle for the active drug, some adverse events occurring at similar frequencies in different treatment groups could be related to lactose vehicle inhalation.

Treatment of Influenza: *Clinical Trials in Adults and Adolescents:* Adverse events that occurred with an incidence $\geq 1.5\%$ in treatment studies are listed in Table 1. This table shows adverse events occurring in patients ≥ 12 years of age receiving RELENZA 10 mg inhaled twice daily, RELENZA in all inhalation regimens, and placebo inhaled twice daily (where placebo consisted of the same lactose vehicle used in RELENZA).

Table 1. Summary of Adverse Events $\geq 1.5\%$ Incidence During Treatment in Adults and Adolescents

| Adverse Event | RELENZA | | Placebo (Lactose Vehicle) (n = 1,520) |
|----------------------------------|---|--|---|
| | 10 mg b.i.d. Inhaled (n = 1,132) | All Dosing Regimens* (n = 2,289) | |
| Body as a whole | | | |
| Headaches | 2% | 2% | 3% |
| Digestive | | | |
| Diarrhea | 3% | 3% | 4% |
| Nausea | 3% | 3% | 3% |
| Vomiting | 1% | 1% | 2% |
| Respiratory | | | |
| Nasal signs and symptoms | 2% | 3% | 3% |
| Bronchitis | 2% | 2% | 3% |
| Cough | 2% | 2% | 3% |
| Sinusitis | 3% | 2% | 2% |
| Ear, nose, and throat infections | 2% | 1% | 2% |
| Nervous system | | | |
| Dizziness | 2% | 1% | <1% |

* Includes studies where RELENZA was administered intranasally (6.4 mg 2 to 4 times per day in addition to inhaled preparation) and/or inhaled more frequently (q.i.d.) than the currently recommended dose.

Additional adverse reactions occurring in less than 1.5% of patients receiving RELENZA included malaise, fatigue, fever, abdominal pain, myalgia, arthralgia, and urticaria.

The most frequent laboratory abnormalities in Phase 3 treatment studies included elevations of liver enzymes and CPK, lymphopenia, and neutropenia. These were reported in similar proportions of zanamivir and lactose vehicle placebo recipients with acute influenza-like illness.

Clinical Trials in Pediatric Patients: Adverse events that occurred with an incidence $\geq 1.5\%$ in children receiving treatment doses of RELENZA in two Phase 3 studies are listed in Table 2. This table shows adverse events occurring in pediatric patients 5 to 12 years old receiving RELENZA 10 mg inhaled twice daily, and placebo inhaled twice daily (where placebo consisted of the same lactose vehicle used in RELENZA).

Table 2. Summary of Adverse Events $\geq 1.5\%$ Incidence During Treatment in Pediatric Patients*

| Adverse Event | RELENZA 10 mg b.i.d. Inhaled (n = 291) | Placebo (Lactose Vehicle) (n = 318) |
|----------------------------------|--|---|
| Respiratory | | |
| Ear, nose, and throat infections | 5% | 5% |
| Ear, nose, and throat hemorrhage | <1% | 2% |
| Asthma | <1% | 2% |
| Cough | <1% | 2% |
| Digestive | | |
| Vomiting | 2% | 3% |
| Diarrhea | 2% | 2% |
| Nausea | <1% | 2% |

* Includes a subset of patients receiving RELENZA for treatment of influenza in a prophylaxis study.

In 1 of the 2 studies described in Table 2, some additional information is available from children (5 to 12 years old) without acute influenza-like illness who received an investigational prophylaxis regimen of RELENZA; 132 children received RELENZA and 145 children received placebo. Among these children, nasal signs and symptoms (zanamivir 20%, placebo 9%), cough (zanamivir 16%, placebo 8%), and throat/tonsil discomfort and pain (zanamivir 11%, placebo 6%) were reported more frequently with RELENZA than placebo. In a subset with chronic pulmonary disease, lower respiratory adverse events (described as asthma, cough, or viral respiratory infections which could include influenza-like symptoms) were reported in 7 of 7 zanamivir recipients and 5 of 12 placebo recipients.

Prophylaxis of Influenza: Family/Household Prophylaxis Studies: Adverse events that occurred with an incidence of $\geq 1.5\%$ in the 2 prophylaxis studies are listed in Table 3. This table shows adverse events occurring in patients ≥ 5 years of age receiving RELENZA 10 mg inhaled once daily for 10 days.

Table 3. Summary of Adverse Events $\geq 1.5\%$ Incidence During 10-Day Prophylaxis Studies in Adults, Adolescents, and Children*

| Adverse Event | Contact Cases | |
|---|------------------------|------------------------|
| | RELENZA (n = 1,068) | Placebo (n = 1,059) |
| Lower respiratory | | |
| Viral respiratory infections | 13% | 19% |
| Cough | 7% | 9% |
| Neurologic | | |
| Headaches | 13% | 14% |
| Ear, nose, and throat | | |
| Nasal signs and symptoms | 12% | 12% |
| Throat and tonsil discomfort and pain | 8% | 9% |
| Nasal inflammation | 1% | 2% |
| Musculoskeletal | | |
| Muscle pain | 3% | 3% |
| Endocrine and metabolic | | |
| Feeding problems (decreased or increased appetite and anorexia) | 2% | 2% |
| Gastrointestinal | | |
| Nausea and vomiting | 1% | 2% |
| Non-site specific | | |
| Malaise and fatigue | 5% | 5% |
| Temperature regulation disturbances (fever and/or chills) | 5% | 4% |

* In prophylaxis studies symptoms associated with influenza-like illness were captured as adverse events; subjects were enrolled during a winter respiratory season during which time any symptoms that occurred were captured as adverse events.

Community Prophylaxis Studies: Adverse events that occurred with an incidence of $\geq 1.5\%$ in 2 prophylaxis studies are listed in Table 4. This table shows adverse events occurring in patients ≥ 5 years of age receiving RELENZA 10 mg inhaled once daily for 28 days.

Table 4. Summary of Adverse Events $\geq 1.5\%$ Incidence During 28-Day Prophylaxis Studies in Adults, Adolescents, and Children*

| Adverse Event | RELENZA (n = 2,231) | Placebo (n = 2,239) |
|---|------------------------|------------------------|
| Neurologic | | |
| Headaches | 24% | 26% |
| Ear, nose, and throat | | |
| Throat and tonsil discomfort and pain | 19% | 20% |
| Nasal signs and symptoms | 12% | 13% |
| Ear, nose, and throat infections | 2% | 2% |
| Lower respiratory | | |
| Cough | 17% | 18% |
| Viral respiratory infections | 3% | 4% |
| Musculoskeletal | | |
| Muscle pain | 8% | 8% |
| Musculoskeletal pain | 6% | 6% |
| Arthralgia and articular rheumatism | 2% | <1% |
| Endocrine and metabolic | | |
| Feeding problems (decreased or increased appetite and anorexia) | 4% | 4% |
| Gastrointestinal | | |
| Nausea and vomiting | 2% | 3% |
| Diarrhea | 2% | 2% |
| Non-site specific | | |
| Temperature regulation disturbances (fever and/or chills) | 9% | 10% |
| Malaise & fatigue | 8% | 8% |

* In prophylaxis studies symptoms associated with influenza-like illness were captured as adverse events; subjects were enrolled during a winter respiratory season during which time any symptoms that occurred were captured as adverse events.

Observed During Clinical Practice: In addition to adverse events reported from clinical trials, the following events have been identified during post-marketing use of zanamivir (RELENZA). Because they are reported voluntarily from a population of unknown size, estimates of frequency cannot be made. These events have been chosen for inclusion due to a combination of their seriousness, frequency of reporting, or potential causal connection to zanamivir (RELENZA).

General: Allergic or allergic-like reaction, including oropharyngeal edema (see PRECAUTIONS).

Cardiac: Arrhythmias, syncope.

Neurologic: Seizures.

Respiratory: Bronchospasm, dyspnea (see WARNINGS and PRECAUTIONS).

Skin: Facial edema; rash, including serious cutaneous reactions (see PRECAUTIONS).

OVERDOSAGE

There have been no reports of overdose from administration of RELENZA. Doses of zanamivir up to 64 mg/day have been administered by nebulizer. Additionally, doses of up to 1,200 mg/day for 5 days have been administered intravenously. Adverse effects were similar to those seen in clinical studies at the recommended dose.

DOSAGE AND ADMINISTRATION

RELENZA is for administration to the respiratory tract by oral inhalation only, using the DISKHALER device provided. **Patients should be instructed in the use of the delivery system. Instructions should include a demonstration whenever possible. If RELENZA is prescribed for children, it should be used only under adult supervision and instruction, and the supervising adult should first be instructed by a healthcare professional (see PRECAUTIONS).**

Patients scheduled to use an inhaled bronchodilator at the same time as RELENZA should use their bronchodilator before taking RELENZA (see WARNINGS and PRECAUTIONS regarding patients with underlying airways disease and other medical conditions).

Treatment: The recommended dose of RELENZA for treatment of influenza in adults and pediatric patients ages 7 years of age and older is 2 inhalations (one 5-mg blister per inhalation for a total dose of 10 mg) twice daily (approximately 12 hours apart) for 5 days. Two doses should be taken on the first day of treatment whenever possible provided there is at least 2 hours between doses. On subsequent days, doses should be about 12 hours apart (e.g., morning and evening) at approximately the same time each day. There are no data on the effectiveness of treatment with RELENZA when initiated more than 2 days after the onset of signs or symptoms.

Prophylaxis: Household Setting: The recommended dose of RELENZA for prophylaxis of influenza in adults and pediatric patients 5 years of age and older in a household setting is 10 mg once daily for 10 days. The 10-mg dose is provided by 2 inhalations (one 5-mg blister per inhalation). The dose should be administered at approximately the same time each day. There are no data on the effectiveness of prophylaxis with RELENZA in a household setting when initiated more than 1.5 days after the onset of signs or symptoms in the index case.

Community Outbreaks: The recommended dose of RELENZA for prophylaxis of influenza in adults and adolescents in a community setting is 10 mg once daily for 28 days. The 10-mg dose is provided by 2 inhalations (one 5-mg blister per inhalation). The dose should be administered at approximately the same time each day. There are no data on the effectiveness of prophylaxis with RELENZA in a community outbreak when initiated more than 5 days after the outbreak was identified in the community. The safety and effectiveness of prophylaxis with RELENZA have not been evaluated for longer than 28 days duration.

HOW SUPPLIED

RELENZA is supplied in a circular double-foil pack (a ROTADISK) containing 4 blisters of the drug. Five ROTADISKS are packaged in a white polypropylene tube. The tube is packaged in a carton with 1 blue and gray DISKHALER inhalation device (NDC 0173-0681-01).

Store at 25°C (77°F); excursions permitted to 15° to 30°C (59° to 86°F) (see USP Controlled Room Temperature). Keep out of reach of children. Do not puncture any RELENZA ROTADISK blister until taking a dose using the DISKHALER.



GlaxoSmithKline
Research Triangle Park, NC 27709

©2006, GlaxoSmithKline. All rights reserved.

March 2006

RL-2270